





0.144	Marie de la	Drotoin	Comme	Chusture	PMC	Taxonomy	OMIM	Books
PubMed Search PubM	Nucleotide	Protein	Genome	Structure	FIVIC	and the same of th	Go Clear	DOOKS
Deuten			ew/Index F	istory (Clipboard	Details Send to Text	, 	
Entrez			eb 18;397(67	20):579-83		R	elated Articl	es, Links
PubMed		nature	999 Feb 18;39			thyloso ooti	vity for n	·CnC
PubMed Services	D	NA.	SK, Ramcha	•		thylase acti Szyf M.	vity 101 ii	СрС
		epartment of mada.	Pharmacolog	y and Thera	peutics, Mo	cGill Universi	ty, Montrea	al, QC,
Related Resources	de de en m is Th	termined by the methylating of codes a methe ethylated cyto translated or the discovery of the methylatery of the discovery o	the enzymation enzyme has not be not	processes on been idealing domain, cytosine, and ansfected in demethylase	of methylat entified: a n bears a der and demethy to human e should pro	ting genome for and demendent and demendent commethylase activates a plasmi mbryonal kide ovide a basis for thylation and	thylation. T mplementa vity that tra id when the ney cells in or the mole	he ry DNA unsforms cDNA vitro. cular
	PN	MID: 100508	51 [PubMed	- indexed fo	r MEDLIN	Œ]		
	Display	Abstract	▼ Show:	20 🔽 Sc	ort [v][Send to Text	नि	

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

0

FILE 'HOME' ENTERED AT 16:55:01 ON 15 MAY 2003)

	FILE 'CAPLUS' ENTERED AT 16:55:38 ON 15 MAY 2003
L1	O S DNA DEMETHYLASE (A) INHIBITOR
L2	29 S DNA DEMETHYLASE
L3	0 S L2 (A) INHIBITION
L4	0 S IMIDAZOLE (A) DNA DEMETHYLASE

2 ANSWER 22 OF 29 CAPLUS COPYRIGHT 2003 ACS AN 1999:218801 CAPLUS DN 131:29241 TI DNA demethylase is a processive enzyme AU Cervoni, Nadia; Bhattacharya, Sanjoy; Szyf, Moshe CS Department of Pharmacology, McGill University, Montreal, QC, H3G 1Y6, Can. Journal of Biological Chemistry (1999), 274(13), 8363-8366 SO CODEN: JBCHA3; ISSN: 0021-9258 PB American Society for Biochemistry and Molecular Biology DTJournal LА English CC 7-4 (Enzymes) DNA methylation patterns are generated during development by a sequence AΒ of methylation and demethylation events. We have recently demonstrated that mammals bear a bona fide demethylase enzyme that removes Me groups from methylated cytosines. A general genome wide demethylation occurs early in development and in differentiating cell lines. This manuscript tests the hypothesis that the demethylase enzyme is a processive enzyme. Using bisulfite mapping, this report demonstrates that demethylase is a processive enzyme and that the rate-limiting step in demethylation is the initiation of demethylation. Initiation of demethylation is detd. by the properties of the sequence. Once initiated, demethylation progresses processively. We suggest that these data provide a mol. explanation for

global hypomethylation.

DNA demethylase hypomethylation

ST

S NCBI	Prot	ein									
PubMed 1	Nucleotide Protein Genome Structure PMC	Taxonomy OMIM Books									
Search Protein	for DNA demethylase	Go Clear									
	Limits Preview/Index History Clipboard	Details									
<u></u>	Display Summary Show; 20 Send to Text										
Entrez Protein											
Linds Floteni	Items 1-20 of 42	Page 1 of 3 Next									
·	☐1: NP_004167 Sterol regulatory element binding transcription factor 1; sterol regulatory element binding protein-1 [Homo sapiens] gi 22547195 ref NP_004167.3 [22547195]										
	Similar to Cytochrome P450 51A1 (CYPL1) (P450L1) (Sterol 14-alpha demethylase) (Lanosterol 14-alpha demethylase) (LDM) (P450-14DM) [Homo sapiens] gi 13650621 ref XP_016472.1 [13650621]										
	☐3: NP_056647 BLink, Domains, L methyl-CpG binding domain protein 2 testis-specific isoform [Homo sapiens] gi 7710145 ref NP_056647.1 [7710145]										
Related resources	☐4: NP_003917 methyl-CpG binding domain protein 3 [Homo sapiengi 4505119 ref NP_003917.1 [4505119]	BLink, Domains, Links									
	□5: NP_003918 BLink, Domains, L methyl-CpG binding domain protein 2 isoform 1 [Homo sapiens] gi 4505117 ref NP_003918.1 [4505117]										
	Cytochrome P450 51 (CYPL1) (P450-L1A1) (Sterol 14-alpha demethylase) (Lanosterol 14-alpha demethylase) (P450-14DM) gi 117176 sp P10614 CP51_YEAST[117176]										
	☐7: P14263 Cytochrome P450 51 (CYPL1) (P450-L1A1) (Sterol : (Lanosterol 14-alpha demethylase) (P450-14DM) gi 117175 sp P14263 CP51_CANTR[117175]	BLink, Domains, Links 14-alpha demethylase)									
	putative cytochrome P450(Obtusifoliol 14-alpha demo (japonica cultivar-group)]	BLink, Domains, Links ethylase) [Oryza sativa									

□9: <u>NP_056723</u>	BLink, Domains, Links
methyl-CpG binding domain protein 1 isoform PCM1 [Homo gi 21464119 ref NP_056723.2 [21464119]	sapiens]
□ 10: NP_056671 methyl-CpG binding domain protein 1 isoform 1 [Homo sapies gi 21464117 ref NP_056671.2 [21464117]	BLink, Domains, Links ns]
□11: NP_056670 methyl-CpG binding domain protein 1 isoform 2 [Homo sapier gi 21464115 ref NP_056670.2 [21464115]	BLink, Domains, Links ns]
□12: NP_604391 cAMP responsive element binding protein 1 isoform B; cAMP element-binding protein-1; active transcription factor CREB; to [Homo sapiens] gi 19745184 ref NP_604391.1 [19745184]	_
□13: NP_002375 methyl-CpG binding domain protein 1 isoform 4 [Homo sapier gi 7710143 ref]NP_002375.1 [7710143]	BLink, Domains, Links ns]
□14: NP_056669 methyl-CpG binding domain protein 1 isoform 3 [Homo sapier gi 7710135 ref NP_056669.1 [7710135]	BLink, Domains, Links ns]
CAMP responsive element binding protein 1 isoform A; cAMP element-binding protein-1; active transcription factor CREB; tr [Homo sapiens] gi 4758054 ref NP_004370.1 [4758054]	-
□16: BAC48664 vanillate O-demethylase oxidoreductase [Bradyrhizobium japo gi 27351658 dbj BAC48664.1 [27351658]	BLink, Domains, Links nicum USDA 110]
□ 17: <u>BAC47655</u> vanillate O-demethylase oxygenase subunit [Bradyrhizobium jagi 27350645 dbj BAC47655.1 [27350645]	BLink, Domains, Links aponicum USDA 110]
18: NP_770039 vanillate O-demethylase oxidoreductase [Bradyrhizobium japogi 27378510 ref NP_770039.1 [27378510]	BLink, Domains, Links nicum]
☐ 19: NP_769030 vanillate O-demethylase oxygenase subunit [Bradyrhizobium jagi 27377501 ref NP_769030.1 [27377501]	BLink, Domains, Links aponicum]

of 3 Next


1

□20: XP_192834 BLink, Domains, Links methyl-CpG binding domain protein 2 [Mus musculus] gi|25054585|ref|XP_192834.1|[25054585] ₹ Display ▼ Show: _20 ▼ Send to Text Summary Page

Items 1-20 of 42

<u>Disclaimer</u> | Write to the Help Desk NCBI | NLM | NIH

0



gi|17554480|ref|NP_498848.1|[17554480]

□29: A31854 BLink, Domains, Links lanosterol 14alpha-demethylase (EC 1.14.14.-) cytochrome P450 51 - yeast (Candida tropicalis) gi|83929|pir||A31854[83929] **□30:** A27491 BLink, Domains, Links lanosterol 14alpha-demethylase (EC 1.14.14.-) cytochrome P450 51 - yeast (Saccharomyces cerevisiae) gi|82967|pir||A27491[82967] □31: Q12645 BLink, Domains, Links Pisatin demethylase (Cytochrome P450 57A1) gi|3929362|sp|Q12645|PID9 FUSSO[3929362] □**32:** BAC20841 BLink, Domains, Links putative cytochrome P450(Obtusifoliol 14-alpha demethylase) [Oryza sativa (japonica cultivar-group)] gi|23617163|dbj|BAC20841.1|[23617163] **☐33:** BAC20835 BLink, Domains, Links putative cytochrome P450(Obtusifoliol 14-alpha demethylase) [Oryza sativa (japonica cultivar-group)] gi|23617157|dbj|BAC20835.1|[23617157] □**34:** NP 524468 BLink, Domains, Links six-banded [Drosophila melanogaster] gi|17738157|ref|NP_524468.1|[17738157] **☐35:** AAH32322 BLink, Domains, Links cytochrome P450, 51 (lanosterol 14-alpha-demethylase) [Homo sapiens] gi|21619563|gb|AAH32322.1|[21619563] **☐36:** AAA53284 BLink, Domains, Links cytochrome P450 lanosterol 14-alpha-demethylase [Candida tropicalis] gi|576547|gb|AAA53284.1|[576547] **□37:** CAB58427 BLink, Links P450 lanosterol 14a demethylase [Cryptococcus neoformans var. neoformans] gi|6065767|emb|CAB58427.1|[6065767] **□38:** <u>BAB53878</u> BLink, Domains, Links probable N-methylproline demethylase [Mesorhizobium loti] gi|14027609|dbi|BAB53878.1|[14027609] **□39:** BAB53454 BLink, Domains, Links vanillate O-demethylase oxidoreductase [Mesorhizobium loti]

0

gi|14026858|dbj|BAB53454.1|[14026858]


□40: BAA15607

BLink, Domains, Links

Vanillate demethylase oxidoreductase (EC 1.-.-.) (Vanillate degradation ferredoxin-like protein). [Escherichia coli] gi|1736436|dbj|BAA15607.1|[1736436]

Display Summ	ary 🔻	Show:	20	T	Send to	Text	₹			
It	ems 21-40	of 42				Previous	Page	2	of 3	Next

<u>Disclaimer</u> | <u>Write to the Help Desk</u> <u>NCBI</u> | <u>NLM</u> | <u>NIH</u>



Related resources

Disclaimer | Write to the Help Desk NCBI | NLM | NIH





PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search Pro	otein 🔻 j	for					Go Clea	ır
	Limits Preview/Inde		dex His	tory	Clipboard	tails		
Display	default ▼	Show: 20	▼ Sen	d to File	▼	Get Subsequence		

BLink, Domains, Links

☐1: NP 056647. methyl-CpG bindin...[gi:7710145] LOCUS MBD2 PRI 06-APR-2003 302 aa linear methyl-CpG binding domain protein 2 testis-specific isoform [Homo DEFINITION sapiens]. ACCESSION NP 056647 VERSION NP 056647.1 GI:7710145 DBSOURCE REFSEQ: accession NM 015832.2 KEYWORDS SOURCE Homo sapiens (human) ORGANISM Homo sapiens Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo. REFERENCE (residues 1 to 302) AUTHORS Krithivas, A., Fujimuro, M., Weidner, M., Young, D.B. and Hayward, S.D. TITLE Protein interactions targeting the latency-associated nuclear antigen of Kaposi's sarcoma-associated herpesvirus to cell JOURNAL J. Virol. 76 (22), 11596-11604 (2002) MEDLINE 22276359 PUBMED 12388720 REMARK GeneRIF: interacts with latency-associated nuclear antigen of Kaposi's Sarcoma-associated herpesvirus (KSHV) to tether KSHV to cell chromosomes REFERENCE 2 (residues 1 to 302) AUTHORS Brackertz, M., Boeke, J., Zhang, R. and Renkawitz, R. TITLE Two highly related p66 proteins comprise a new family of potent transcriptional repressors interacting with MBD2 and MBD3 JOURNAL J. Biol. Chem. 277 (43), 40958-40966 (2002) MEDLINE 22287387 PUBMED 12183469 REMARK GeneRIF: interaction with two highly related p66 proteins REFERENCE (residues 1 to 302) AUTHORS Detich, N., Theberge, J. and Szyf, M. TITLE Promoter-specific activation and demethylation by MBD2/demethylase J. Biol. Chem. 277 (39), 35791-35794 (2002) JOURNAL MEDLINE 22229441 PUBMED 12177048 REMARK GeneRIF: MBD2 protein activates CpG sites within the promoter region of reporter genes REFERENCE (residues 1 to 302) 4 Bakker, J., Lin, X. and Nelson, W.G. AUTHORS

TITLE Methyl-CpG binding domain protein 2 represses transcription from hypermethylated pi-class glutathione S-transferase gene promoters in hepatocellular carcinoma cells

JOURNAL J. Biol. Chem. 277 (25), 22573-22580 (2002)

MEDIATNE 22063361 PUBMED 11960994

REMARK GeneRIF: Methyl-CpG binding domain protein 2 represses transcription from hypermethylated pi-class glutathione

S-transferase gene promoters in hepatocellular carcinoma cells

REFERENCE (residues 1 to 302)

AUTHORS Ng, H.H., Zhang, Y., Hendrich, B., Johnson, C.A., Turner, B.M., Erdjument-Bromage, H., Tempst, P., Reinberg, D. and Bird, A.

```
TITLE
            MBD2 is a transcriptional repressor belonging to the MeCP1 histone
             deacetylase complex
  JOURNAL
            Nat. Genet. 23 (1), 58-61 (1999)
  MEDLINE
             99400553
   PUBMED
            10471499
REFERENCE
                (residues 1 to 302)
            6
  AUTHORS
            Hendrich, B., Abbott, C., McQueen, H., Chambers, D., Cross, S. and
  TITLE
            Genomic structure and chromosomal mapping of the murine and human
            Mbd1, Mbd2, Mbd3, and Mbd4 genes
  JOURNAL
            Mamm. Genome 10 (9), 906-912 (1999)
  MEDLINE
            99373255
            10441743
   PUBMED
REFERENCE
            7
                (residues 1 to 302)
  AUTHORS
            Bhattacharya, S.K., Ramchandani, S., Cervoni, N. and Szyf, M.
  TITLE
            A mammalian protein with specific demethylase activity for mCpG DNA
  JOURNAL
            Nature 397 (6720), 579-583 (1999)
  MEDLINE
            99158138
   PUBMED
            10050851
REFERENCE
            8
                (residues 1 to 302)
  AUTHORS
            Hendrich, B. and Bird, A.
  TITLE
            Identification and characterization of a family of mammalian
            methyl-CpG binding proteins
  JOURNAL
            Mol. Cell. Biol. 18 (11), 6538-6547 (1998)
  MEDLINE
            98449942
   PUBMED
            9774669
REFERENCE
            9
                (residues 1 to 302)
  AUTHORS
            Scanlan, M.J., Chen, Y.T., Williamson, B., Gure, A.O., Stockert, E.,
            Gordan, J.D., Tureci, O., Sahin, U., Pfreundschuh, M. and Old, L.J.
  TITLE
            Characterization of human colon cancer antigens recognized by
            autologous antibodies
  JOURNAL
            Int. J. Cancer 76 (5), 652-658 (1998)
  MEDLINE
            98272252
   PUBMED
            9610721
COMMENT
            REVIEWED REFSEQ: This record has been curated by NCBI staff. The
            reference sequence was derived from AF120989.1, AF120989.1 and
            AF120994.1.
```

Summary: DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of binding specifically to methylated DNA. MECP2, MBD1 and MBD2 can also repress transcription from methylated gene promoters. MBD2 may function as mediators of the biological consequences of the methylation signal. It is also reported that the MBD2 protein functions as a demethylase to activate transcription, as DNA methylation causes gene silencing. However, MBD2 in HeLa cells does not demethylate DNA, probably due to HeLa cell's using an alternative pathway involving MBD2 to silence methylated genes.

Transcript Variant: This variant (testis-specific) includes an alternate exon located within intron 2 resulting in a distinct COOH terminus.

```
FEATURES
```

Location/Qualifiers

source

1..302

/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="18"

/map="18q21" 1..302

Protein

/product="methyl-CpG binding domain protein 2

testis-specific isoform"

```
Region
                            /region_name="DNA binding domain found in several
                            chromosomal proteins including the Methyl CpG binding
                            /note="tam"
                            /db_xref="CDD:LOAD tam"
      CDS
                            1..\overline{302}
                            /gene="MBD2"
                            /coded_by="NM_015832.2:230..1138"
                            /db_xref="LocusID:8932"
                            /db_xref="MIM:603547"
ORIGIN
          1 mrahpgggrc cpeqeegesa aggsgaggds aieqggqgsa lapspvsgvr regargggrg
         61 rgrwkqagrg ggvcgrgrgr grgrgrgrgr grgrgrppsg gsglggdggg cggggsgggg
       121 aprrepvpfp sgsagpgprg pratesgkrm dcpalppgwk keevirksgl sagksdvyyf
181 spsgkkfrsk pqlarylgnt vdlssfdfrt gkmmpsklqk nkqrlrndpl nqnklrwnth
241 rpapwhalsr lcllircllc lecayplplh lvnsyssktq lhclhlweac paysrqnqsf
       301 pp
//
```

<u>Disclaimer</u> | <u>Write to the Help Desk</u> <u>NCBI</u> | <u>NLM</u> | <u>NIH</u>





PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search Pro	otein 🔻 fo	or					Go Clea	ar
Limits		Preview/Index History		tory	Clipboard	De	Details	
Display	default ▼	Show: 20	▼ Send	to File	▼	Get Subsequence		

BLink, Domains, Links

☐1: NP_056647. methyl-CpG bindin...[gi:7710145] LOCUS MBD2 302 aa linear PRI 06-APR-2003 DEFINITION methyl-CpG binding domain protein 2 testis-specific isoform [Homo sapiens]. ACCESSION NP 056647 VERSION NP 056647.1 GI:7710145 DBSOURCE REFSEQ: accession NM 015832.2 KEYWORDS SOURCE Homo sapiens (human) ORGANISM Homo sapiens Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo. REFERENCE (residues 1 to 302) AUTHORS Krithivas, A., Fujimuro, M., Weidner, M., Young, D.B. and Hayward, S.D. TITLE Protein interactions targeting the latency-associated nuclear antigen of Kaposi's sarcoma-associated herpesvirus to cell chromosomes JOURNAL J. Virol. 76 (22), 11596-11604 (2002) MEDLINE 22276359 PUBMED 12388720 REMARK GeneRIF: interacts with latency-associated nuclear antigen of Kaposi's Sarcoma-associated herpesvirus (KSHV) to tether KSHV to cell chromosomes REFERENCE 2 (residues 1 to 302) Brackertz, M., Boeke, J., Zhang, R. and Renkawitz, R. AUTHORS TITLE Two highly related p66 proteins comprise a new family of potent transcriptional repressors interacting with MBD2 and MBD3 JOURNAL J. Biol. Chem. 277 (43), 40958-40966 (2002) MEDLINE 22287387 PUBMED 12183469 GeneRIF: interaction with two highly related p66 proteins REMARK REFERENCE (residues 1 to 302) AUTHORS Detich, N., Theberge, J. and Szyf, M. TITLE Promoter-specific activation and demethylation by MBD2/demethylase JOURNAL J. Biol. Chem. 277 (39), 35791-35794 (2002) MEDLINE 22229441 PUBMED 12177048 REMARK GeneRIF: MBD2 protein activates CpG sites within the promoter

region of reporter genes

(residues 1 to 302) REFERENCE

AUTHORS Bakker, J., Lin, X. and Nelson, W.G.

TITLE Methyl-CpG binding domain protein 2 represses transcription from hypermethylated pi-class glutathione S-transferase gene promoters in hepatocellular carcinoma cells

JOURNAL J. Biol. Chem. 277 (25), 22573-22580 (2002)

22063361 MEDLINE PUBMED 11960994

REMARK GeneRIF: Methyl-CpG binding domain protein 2 represses transcription from hypermethylated pi-class glutathione

S-transferase gene promoters in hepatocellular carcinoma cells

REFERENCE (residues 1 to 302)

AUTHORS Ng, H.H., Zhang, Y., Hendrich, B., Johnson, C.A., Turner, B.M., Erdjument-Bromage, H., Tempst, P., Reinberg, D. and Bird, A.

```
MBD2 is a transcriptional repressor belonging to the MeCP1 histone
   TITLE
            deacetylase complex
  JOURNAL
            Nat. Genet. 23 (1), 58-61 (1999)
  MEDLINE
            99400553
            10471499
   PUBMED
REFERENCE
            6
               (residues 1 to 302)
  AUTHORS
            Hendrich, B., Abbott, C., McQueen, H., Chambers, D., Cross, S. and
  TITLE
            Genomic structure and chromosomal mapping of the murine and human
            Mbd1, Mbd2, Mbd3, and Mbd4 genes
  JOURNAL
            Mamm. Genome 10 (9), 906-912 (1999)
  MEDLINE
            99373255
   PUBMED
            10441743
REFERENCE
            7
               (residues 1 to 302)
  AUTHORS
            Bhattacharya, S.K., Ramchandani, S., Cervoni, N. and Szyf, M.
  TITLE
            A mammalian protein with specific demethylase activity for mCpG DNA
  JOURNAL
            Nature 397 (6720), 579-583 (1999)
  MEDLINE
            99158138
   PUBMED
            10050851
REFERENCE
            8 (residues 1 to 302)
  AUTHORS
            Hendrich, B. and Bird, A.
  TITLE
            Identification and characterization of a family of mammalian
            methyl-CpG binding proteins
  JOURNAL
            Mol. Cell. Biol. 18 (11), 6538-6547 (1998)
  MEDLINE
            98449942
   PUBMED
            9774669
REFERENCE
               (residues 1 to 302)
  AUTHORS
            Scanlan, M.J., Chen, Y.T., Williamson, B., Gure, A.O., Stockert, E.,
            Gordan, J.D., Tureci, O., Sahin, U., Pfreundschuh, M. and Old, L.J.
            Characterization of human colon cancer antigens recognized by
  TITLE
            autologous antibodies
 JOURNAL
            Int. J. Cancer 76 (5), 652-658 (1998)
 MEDLINE
            98272252
   PUBMED
            9610721
COMMENT
            REVIEWED REFSEQ: This record has been curated by NCBI staff. The
            reference sequence was derived from AF120989.1, AF120989.1 and
            AF120994.1.
```

Summary: DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of binding specifically to methylated DNA. MECP2, MBD1 and MBD2 can also repress transcription from methylated gene promoters. MBD2 may function as mediators of the biological consequences of the methylation signal. It is also reported that the MBD2 protein functions as a demethylase to activate transcription, as DNA methylation causes gene silencing. However, MBD2 in HeLa cells does not demethylate DNA, probably due to HeLa cell's using an alternative pathway involving MBD2 to silence methylated genes.

Transcript Variant: This variant (testis-specific) includes an alternate exon located within intron 2 resulting in a distinct COOH terminus.

```
Location/Qualifiers
source
                1..302
                /organism="Homo sapiens"
                /db_xref="taxon:9606"
                /chromosome="18"
                /map="18g21"
Protein
                1..302
                /product="methyl-CpG binding domain protein 2
                testis-specific isoform"
```

FEATURES

```
Region
                            147..217
                            /region name="DNA binding domain found in several
                            chromosomal proteins including the Methyl CpG binding
                            /note="tam"
                            /db_xref="CDD:LOAD_tam"
      CDS
                            1..\overline{3}02
                            /gene="MBD2"
                            /coded_by="NM_015832.2:230..1138"
                            /db xref="LocusID:8932"
                            /db_xref="MIM:603547"
ORIGIN
           1 mrahpgggrc cpeqeegesa aggsgaggds aleqggqgsa lapspvsgvr regargggrg
        61 rgrwkqagrg ggvcgrgrgr grgrgrgrgr grgrgrppsg gsglggdggg cggggsgggg
121 aprrepvpfp sgsagpgprg pratesgkrm dcpalppgwk keevirksgl sagksdvyyf
        181 spsgkkfrsk pqlarylgnt vdlssfdfrt gkmmpsklqk nkqrlrndpl nqnklrwnth
241 rpapwhalsr lcllircllc lecayplplh lvnsyssktq lhclhlweac paysrqnqsf
//
```

<u>Disclaimer | Write to the Help Desk</u> <u>NCBI | NLM | NIH</u>







PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonom	y OMIM	Books			
Search PubMe	d ▼ for						Go Clea	r			
	Limits	Previe	w/Index H	istory Cl	lipboard	Details					
)			<u>-</u>							
Č	Display Ab	stract	▼ Show:	20 ▼ Sort	▼	Send to Tex	d ▼				
Entrez	□1: Natur	□1: Nature 1999 Feb 18;397(6720):568-9									
PubMed	Comm	ent on:									
			99 Feb 18;39°	7(6720):579-	<u>83.</u>						
		ature									
	Gene	express	ion. The a	nazing der	nethyla	se.					
PubMed Services	Cedar	H, Verdi	ine GL.	•							
	Publica	ation Type	es:								
		omment									
	• N	ews									
	PMID:	1005084	7 [PubMed -	indexed for l	MEDLIN	E]					
Related Resources	Display Ab	stract	▼ Show:	20 ▼ Sort	▼[:	Send to Tex	t 🔻				

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

.







PubM	ed	Nucleotide	Protein	Genome	Structu	re PMC	C Taxo	nomy	OMIM	Books	
Search	PubMed	▼ for							Go Clear		
		Limits	Previe	w/Index H	istory	Clipboar	d De	tails	•		
				···— <u>—</u> ·····	*****						
		Display A	bstract	▼ Show:	20 ▼	Sort ▼	Send to	Text	▼		
		□1: Proc	Natl Acad	Sci U S A 19	999 May	25:96(11):	5804-6	De	loted Autic	مادا اما	
Entrez			- 1-01		ood iviay	23,70(11).	.2024-0	Re	lated Artic	ies, Links	
PubMed			nent on:								
		0 <u>F</u>	Proc Natl A	cad Sci U S	A. 1999 I	May 25;96	(11):6107	<u>-12.</u>			
			w.pnas.org	in PubMed C							
		DNA	demeth	ylation.							
PubMed											
Services	;	Wolff	e AP, Jon	es PL, Wade	PA.						
		Laboratory of Molecular Embryology, National Institute of Child Health and Human									
		Development, National Institutes of Health, Bethesda, MD 20892-5431, USA.									
		Public	ation Type	es:							
		。 C	Comment								
			eview						,		
		• R	eview, Tu	torial					(
Related Resource	∋s	PMID	: 10339513	3 [PubMed -	indexed for	or MEDLI	NE]				
		Display Ab	stract	▼ Show: 3	n						
		Display AU	ou a Cl	▼ Show: _2	20 ▼ S	ort ▼	Send to	Text	₹		

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

0